


Amendments to the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

Claims 1-4 (withdrawn)

5. (Previously presented) A method of treating, ameliorating, or preventing seizures associated with epilepsy in a subject in need thereof, the method comprising:

administering a pharmaceutical composition comprising about 0.1 to about 500 mg of agmatine or an agmatine analog, or a pharmaceutically acceptable salt thereof per kilogram of the subject's weight to treat, reduce, or prevent seizures associated with epilepsy in the subject, wherein the agmatine analog has the following formula



wherein n is 0 to about 10;

R₁, R₂, R₃, R₄, and R₅, are each independently, or any combination thereof: hydrogen, hydroxy, substituted or unsubstituted C₁₋₁₀ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted arylalkyl (comprising Ar-(CH₂)_m; where Ar is aromatic and m is 0 to about 10) substituted or unsubstituted C₁₋₁₀ alkoxy, substituted or unsubstituted C₁₋₁₀ acyl, halogeno, amido, phenyl, thio, or amino; and

X and Y are each independently: O, NH, CH₂, CF₂, Se, C=O, C=N, or C=S, or X-Y together is HC=CH, C≡C, N=N, N=CH, CH=N, or a saturated or unsaturated ring.

6. (Canceled)

7. (Original) A method according to claim 5, wherein the pharmaceutical composition comprises agmatine or its pharmaceutically acceptable salt and a pharmaceutically acceptable carrier.

8. (Canceled).

9. (Currently amended) A method according to claim 87, wherein the composition is administered in a dose of about 0.1 to about 50 mg/kg per day indefinitely or until seizures associated with the epilepsy.

10. (Canceled).


11. (Previously presented) A method according to claim 5, comprising preventing or reducing seizure activity.

12. (Canceled).

13. (Previously presented) A method of treating or preventing seizures associated with epilepsy in a human comprising:

identifying a human subject in need of said treatment or prevention; and

administering about 0.1 to about 500 mg of agmatine or an agmatine analog, or a pharmaceutically acceptable salt thereof per kilogram of the subject's weight to the human subject, wherein the agmatine analog has the following formula



wherein n is 0 to about 10;

R₁, R₂, R₃, R₄, and R₅, are each independently, or any combination thereof: hydrogen, hydroxy, substituted or unsubstituted C₁₋₁₀ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted arylalkyl (comprising Ar-(CH₂)_m; where Ar is aromatic and m is 0 to about 10) substituted or unsubstituted C₁₋₁₀ alkoxy, substituted or unsubstituted C₁₋₁₀ acyl, halogeno, amido, phenyl, thio, or amino; and

X and Y are each independently: O, NH, CH₂, CF₂, Se, C=O, C=N, or C=S, or X-Y together is HC=CH, C≡C, N=N, N=CH, CH=N, or a saturated or unsaturated ring.

14. (Original) A method according to claim 13, comprising identifying a human subject in need of said treatment by analyzing an electroencephalogram taken of the human subject.

15. (Previously presented) A method according to claim 13, comprising identifying a human subject in need of said treatment by observing the occurrence of a seizure in said subject.

16. (Previously presented). A method according to claim 13, comprising administering the effective amount of agmatine, an agmatine analog or a pharmaceutically acceptable salt thereof to the human subject indefinitely or until the seizures associated with epilepsy cease.

17. (Previously presented) A method according to claim 13, comprising preventing or reducing seizures associated with epileptic activity.

18. (Original) A method according to claim 13, comprising administering the effective amount of agmatine, an agmatine analog or a pharmaceutically acceptable salt thereof as a pharmaceutical composition.

19. (Original) A method according to claim 13, comprising administering the effective amount of agmatine, an agmatine analog or a pharmaceutically acceptable salt thereof parenterally.

Application No.: 09/881,215

20. (Original). A method according to claim 13, comprising administering the effective amount of agmatine, an agmatine analog or a pharmaceutically acceptable salt thereof orally.